

AMCoR

Asahikawa Medical University Repository <http://amcor.asahikawa-med.ac.jp/>

Hypertension Research (2008.02) 31巻2号:193~201.

A Newly Estimated Glomerular Filtration Rate Is Independently
Associated with Arterial Stiffness in Japanese Patients
(新たに提案された推定糸球体濾過値は日本人患者の動脈硬度と独立に
関連する)

Nakagawa Naoki, Takahashi Fumihiko, Chinda Junko,
Kobayashi Motoi, Hayashi Yoshikazu, Abe Masahiko, Saijo
Yasuaki, Kikuchi Kenjiro, Hasebe Naoyuki

A Newly Estimated Glomerular Filtration Rate Is Independently Associated with Arterial Stiffness in Japanese Patients

Naoki Nakagawa^{1,2}, Fumihiko Takahashi², Junko Chinda², Motoi Kobayashi²,
Yoshikazu Hayashi², Masahiko Abe², Yasuaki Saijo³,
Kenjiro Kikuchi¹ and Naoyuki Hasebe¹

¹ Cardiovascular Division, Department of Internal Medicine and ³ Department of Health Science, Asahikawa Medical College, 2-1-1-1 Midorigaoka-higashi, Asahikawa 078-8510, Japan

² Department of Internal Medicine, Hokkaido Prefectural Haboro Hospital, 110 Sakae, Haboro, 078-4197, Japan

Running title: Estimated GFR and Arterial Stiffness

Corresponding author: Naoki Nakagawa, M.D., Cardiovascular Division, Department of Internal Medicine, Asahikawa Medical College

Phone : +81-166-68-2442, FAX : +81-166-68-2449

E-mail : naka-nao@asahikawa-med.ac.jp

Total pages: 21

Figures: 4

Tables: 5

Abstract

Chronic kidney disease (CKD) is associated with an increased risk of cardiovascular disease, and thus is a major worldwide public health problem. Recently, an estimated glomerular filtration rate (eGFR) using the Modification of Diet in Renal Disease equation for Japanese patients was proposed by the Japanese Society of Nephrology. However, the role of eGFR in the assessment of atherosclerosis is not well understood in Japanese patients. We analyzed the relationship between eGFR and severity of arterial stiffness using brachial-ankle pulse wave velocity (baPWV) in 647 adult Japanese patients. baPWV correlated significantly and positively with age, hypertension, diabetes, prior cardiovascular disease, blood pressure, pulse pressure and heart rate, and negatively with eGFR ($r=-0.405$, $p<0.001$). A multiple regression analysis revealed that baPWV correlated independently with eGFR. Furthermore, there was a stepwise increase in baPWV, corresponding to advances in CKD through stages 1 to 5. When CKD stage 3 was divided at eGFR 45mL/min/1.73m², the baPWV of stage 3b (eGFR 30 to 44) was significantly higher than that of stage 3a (eGFR 45 to 59) independent of traditional risk factors, suggesting that an eGFR of 45mL/min/1.73m² may be a critical cut off value to predict arterial stiffness in CKD. In conclusion, the newly proposed eGFR is significantly associated with arterial stiffness, independent of traditional risk factors for cardiovascular disease.

Key words: chronic kidney disease, estimated glomerular filtration rate, arterial stiffness, cross-sectional studies, Japanese patients.

Introduction

Chronic kidney disease (CKD) is a major worldwide public health problem, with adverse outcomes of kidney failure, cardiovascular disease (CVD), and premature death (1-3). A number of prospective epidemiologic studies have shown that patients with CKD are at increased risk for CVD, independent of conventional cardiovascular risk factors (4-6). Thus, the National Kidney Foundation formed a task force to heighten awareness of CVD in CKD (7), and defined CKD as either (1) kidney damage for ≥ 3 months, confirmed by kidney biopsy or markers of kidney damage, with or without a decrease in glomerular filtration rate (GFR), or (2) GFR <60 mL/min/1.73m² for ≥ 3 months, with or without kidney damage (8). Pulse wave velocity (PWV) is known to be an indicator of arterial stiffness (9, 10), and an independent predictor of cardiovascular events in patients with end-stage renal disease (11, 12). A simple and noninvasive method to automatically measure brachial-ankle PWV (baPWV) was recently developed to screen large populations (13, 14). In several reports that included patients with different stages of CKD, PWV increased proportionally to decreased eGFR or creatinine clearance (15-17). Recently, an estimated GFR (eGFR) for Japanese patients was proposed by the Japanese Society of Nephrology (18); however, its clinical usefulness has not yet been fully established. In this study, we investigated for the first time the significance of eGFR as a predictor of arterial stiffness in Japanese patients to clarify whether CKD is related to the progression of arterial stiffness in such patients.

METHODS

Subjects

Subjects were 647 consecutive patients (318 men and 329 women; aged 15 to 92 years), who underwent brachial-ankle pulse wave velocity (baPWV) at Hokkaido Prefectural Haboro Hospital from January 2005 to April 2006. Patients on hemodialysis or with an ankle brachial pressure index (ABI) of ≤ 0.9 were excluded. Age, gender, lipid parameters, and conventional cardiovascular risk factors were recorded. Body weight and height were measured during the examination in light indoor clothing without shoes. baPWV, ABI, heart rate (HR), and blood pressure (BP) were measured with a pulse pressure analyzer (model: BP-203RPE II; Nihon Colin, Tokyo, Japan) as described previously (13, 14). PWV was expressed in centimeters per second. Fasting or non-fasting blood samples were drawn from the antecubital vein of seated participants with minimal tourniquet use. Samples were collected into vacuum tubes containing ethylenediaminetetraacetic acid for hemoglobin (Hb) and HbA1c measurement, or a serum separator gel for serum creatinine, total cholesterol (TC), low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, triglycerides, and uric acid measurement. A history of smoking was defined as > 10 pack-years. Hypertension was defined as either systolic pressure ≥ 140 and/or diastolic pressure ≥ 90 mmHg, or current use of anti-hypertensive medications. Diabetes was defined as one of the following: fasting blood sugar ≥ 126 mg/dl; non-fasting blood sugar ≥ 200 mg/dl or HbA1c $\geq 6.5\%$; or current use of insulin or an oral hypoglycemic agent. Dyslipidemia was defined as: TC ≥ 220 mg/dl; HDL cholesterol ≤ 40 mg/dl; triglycerides ≥ 150 mg/dl; or current use of an

anti-hyperlipidemic medication. The dipstick urinalysis (Bayer, Tokyo, Japan) for proteinuria was performed on spontaneously voided fresh urine. Urine test results were interpreted by the physicians and recorded as (-), (\pm), 1+, 2+, and 3+. Results recorded as (-) and (\pm) are defined as the absence of proteinuria; others are defined as the presence of proteinuria (7.3% of total subjects). Body mass index (BMI) was calculated as weight divided by height squared.

This study was conducted in accordance with the guidelines of the ethics committee of Hokkaido Prefectural Haboro hospital. Informed consent was obtained from each patient before entry into the study.

Measurement of Estimated Glomerular Filtration Rate

Glomerular filtration rate (GFR) was estimated from the MDRD equation for Japanese patients, recently proposed by a working group of the Japanese Chronic Kidney Disease Initiative (18):

Estimated GFR (eGFR) = $186.3 \times (\text{serum creatinine})^{-1.154} \times \text{age}^{-0.203} \times (0.742 \text{ if female}) \times (0.881 \text{ if Japanese})$.

Serum creatinine was analyzed by the enzyme method (normal: 0.7–1.2 mg/dl for men, 0.5–0.8 mg/dl for women) using a Toshiba TBA-200FR automated analyzer (Toshiba, Tokyo, Japan). Serum creatinine was calibrated using the following formula: serum creatinine (Yaffe method) = 0.2 + serum creatinine (enzyme method) (19, 20). eGFR was analyzed using the following five categories provided by the Kidney Disease Outcomes Quality Initiative (K/DOQI) (8): stage 1: $\text{GFR} \geq 90 \text{ mL/min/1.73m}^2$; stage 2: $\text{GFR} 60 \text{ to } 89$

mL/min/1.73m²; stage 3: GFR 30 to 59 mL/min/1.73m²; stage 4: GFR 15 to 30 mL/min/1.73m²; and stage 5: GFR <15 mL/min/1.73m². Furthermore, we calculated estimated creatinine clearance (eCcr) using the Cockcroft-Gault equation, which is often used clinically, and examined correlations between eCcr and baPWV, and between eCcr and eGFR.

Statistical analysis

Results were expressed as the means±SD or number (percentage). Univariate and multivariate linear regression was used for continuous variables. The unadjusted mean of baPWV was compared to the different CKD stages using eGFR, with one-way analysis of variance (ANOVA) followed by multiple comparisons using Tukey's Studentized Range Test. Furthermore, additional comparisons were made under the condition that category CKD 3 was divided into early stage 3a (GFR 45 to 59 mL/min/1.73m²) and late stage 3b (GFR 30 to 44 mL/min/1.73m²) as previously reported (6). The adjusted mean of baPWV was compared among the different CKD stages, with analysis of covariance (ANCOVA) for age, sex, systolic BP, BMI, HR, Hb, TC, prior CVD, smoking, medication for hypertension, medication for diabetes, and medication for dyslipidemia. For the analysis of the amount of proteinuria at each CKD stage, proteinuria was scored as follows: dipstick findings of (-) and (±) were scored as 0, and findings of (1+), (2+), and (3+) were scored as 1, 2, and 3, respectively. *p* values <0.05 were considered statistically significant. All statistical analyses were performed with the SPSS software package Version 11.0 for Windows (SPSS Inc., Chicago, IL).

RESULTS

The baseline characteristics of the subjects are outlined in Table 1. The mean age was 66.3 ± 11.7 years (range, 15 to 92 years). The prevalence of hypertension and diabetes mellitus were about 80% and 35%, respectively. The mean body mass index (BMI) was 25.9 kg/m^2 . The mean serum creatinine was $0.84 \pm 0.61 \text{ mg/dl}$ (men, $0.97 \pm 0.79 \text{ mg/dl}$; women, $0.72 \pm 0.33 \text{ mg/dl}$). The mean eGFR using the MDRD equation for Japanese patients was 63.4 ± 14.5 (men, 66.1 ± 15.8 ; women, $60.9 \pm 12.7 \text{ mL/min/1.73m}^2$). The mean baPWV was $1775 \pm 375 \text{ cm/s}$ (range, 950 to 3341), $1782 \pm 376 \text{ cm/s}$ for men, and $1768 \pm 374 \text{ cm/s}$ for women. Univariate correlations between baPWV and various parameters are listed in Table 2. baPWV correlated significantly and positively with age, hypertension, diabetes, prior CVD, systolic, diastolic and mean BP, pulse pressure, HR, and serum creatinine, and negatively with eGFR ($r=-0.405$, $p<0.0001$), eCcr, Hb, BMI, and TC in the univariate linear regression analysis. However, baPWV did not significantly correlate with sex, smoking, triglycerides, HDL cholesterol, LDL cholesterol, uric acid (in a subset of 607 subjects for whom triglycerides, HDL and LDL cholesterol data were available), or HbA1c (in a subset of 278 subjects for whom HbA1c data was available). Figure 1 shows the scatter chart of the relationship between baPWV and eGFR. A similar negative relationship between eGFR and baPWV was observed in both men and women (men, $r=-0.441$; women, $r=-0.390$; both $p<0.0001$) and in both diabetes and non-diabetes (diabetes, $r=-0.406$; non-diabetes, $r=-0.400$; both $p<0.0001$). Furthermore, eCcr calculated using the Cockcroft-Gault equation was associated significantly and negatively with baPWV ($r=-0.555$, $p<0.0001$) and positively with eGFR ($r=0.705$, $p<0.0001$).

Although eCcr was also negatively associated with baPWV, we performed analyses using eGFR by the MDRD study equation modified for Japanese patients because eGFR was a more accurate predictor of GFR than eCcr (18). The calculation of eGFR itself includes age, and baPWV per se is strongly influenced by age. To exclude possible confounding problems of age, we performed a multiple regression analysis for baPWV. The multiple regression analysis revealed that baPWV was correlated independently with eGFR ($r=-0.125$, $p<0.0001$), age ($r=0.498$, $p<0.0001$), systolic BP ($r=0.374$, $p<0.0001$), HR ($r=0.220$, $p<0.0001$), BMI ($r=-0.148$, $p<0.0001$) and diabetes (diabetes=1, non-diabetes=0) ($r=0.058$, $p=0.042$) (Table 3), suggesting that eGFR was negatively correlated with baPWV independent of traditional risk factors.

Next, we investigated the relationship between baPWV and the different CKD stages according to the K/DOQI category of kidney function. Figure 2 shows the distribution of eGFR and successive categories of kidney function (stage 1: GFR ≥ 90 ; stage 2: GFR 60 to 89; stage 3: GFR 30 to 59; stage 4: GFR 15 to 30; and stage 5: GFR <15 L/min/1.73m²). The average creatinine was 0.57 mg/dl, 0.70 mg/dl, 0.93 mg/dl, 2.16 mg/dl and 5.92 mg/dl in stages 1-5, respectively. eGFR distribution was shifted toward lower values in women compared to men. The prevalences of CKD stages 1-5 were 2.6, 60.6, 34.2, 1.9 and 0.8%, representing 17, 392, 221, 12 and 5 patients, respectively. Overall, the prevalence of eGFR <60 mL/min/1.73m² was 36.9 %, representing 238 patients (28.3% or 90 men and 45.0% or 148 women). Furthermore, there was a stepwise increase in baPWVs, corresponding to advances in CKD through stages 1 to 5 ($p<0.0001$ for trend) (Figure 3A). After adjustment for age, sex, systolic BP, BMI, HR, Hb, TC, prior CVD, smoking,

medication for hypertension, medication for diabetes, and medication for dyslipidemia, significant differences in baPWV were observed across the different CKD stages ($p<0.0001$) (Figure 3B). When CKD stage 3, moderately reduced GFR, was divided at eGFR 45 mL/min/1.73m² (stage 3a: eGFR 45 to 59; stage 3b: eGFR 30 to 44 mL/min/1.73m²), the adjusted baPWV in stage 3b increased significantly compared to that in stage 3a (1783cm/sec versus 1889cm/sec, $p<0.0001$) (Figure 3C) independent of age, systolic BP and other traditional risk factors. Table 4 shows the differences in parameters at each CKD stage. Significant differences in age, systolic BP, eGFR, baPWV, and the prevalence of hypertension, diabetes, dyslipidemia, and prior CVD were observed across the CKD stages. In addition, we also investigated the relationship between the prevalence of proteinuria and baPWV in a subset of 589 subjects for whom dipstick findings for proteinuria were available. Table 5 lists the ages, the systolic BP, eGFR, and baPWV values, and the prevalence of hypertension, diabetes mellitus, dyslipidemia, and prior CVD according to dipstick proteinuria findings. Higher proteinuria levels were associated with greater systolic BP, lower eGFR, greater baPWV, and greater prevalence of diabetes mellitus and dyslipidemia. The proteinuria score also increased with the progression of CKD stage ($p<0.0001$ for trend) and was significantly different between CKD stages 3a and 3b ($p<0.001$) (Figure 4), suggesting that an eGFR of 45 mL/min/1.73m² may be a critical cut off value not only to predict arterial stiffness progression, but also to predict the development of end-stage renal disease in CKD.

DISCUSSION

Our study demonstrated that increased baPWV is associated with decreased eGFR and advanced CKD stages, independent of traditional cardiovascular risk factors such as age, smoking, BMI, systolic blood pressure, prior CVD, hypertension, diabetes and dyslipidemia, in patients living in rural Japan. Furthermore, to our knowledge, this is the first report to demonstrate an association between arterial stiffness and eGFR calculated using the MDRD equation for Japanese patients.

In the present study, we calculated eGFR using the MDRD equation, as proposed by the Japanese Society of Nephrology, which considers race in the calculation and has been adapted for Japanese subjects. Although the equation is now being assessed for validity, the Japanese Society of Nephrology has indicated that there is a good correlation between eGFR—particularly low eGFR (<60 mL/min/1.73m²)—and the actual GFR measured by inulin clearance, the gold standard for measuring GFR, in Japanese patients (18). Meanwhile, the Cockcroft-Gault equation is often used to estimate GFR in clinical settings, but this equation is originally estimated creatinine clearance, not inulin clearance which is a gold standard as measured GFR. Several studies have shown that the Cockcroft-Gault equation overestimates GFR, especially in patients with moderate and severe CKD, which was defined as $GFR < 60$ mL/min/1.73m² in the MDRD study (18, 21). Furthermore, a recent advisory from the American Heart Association Kidney and Cardiovascular Disease Council formed in collaboration with the National Kidney Foundation revealed that CKD can be reliably detected with the combined use of the MDRD equation for eGFR and a sensitivity test to detect microalbuminuria (22). Thus, we used the MDRD equation for

Japanese patients, which is suitable for routine clinical use or large screening programs for the detection of early stages of CKD.

Firstly, we demonstrated that age, hypertension (i.e., systolic, diastolic and mean BP and pulse pressure), diabetes mellitus, and prior CVD had good positive correlations with baPWV, and that eGFR for Japanese patients had both a significant and negative association with baPWV, independent of other cardiovascular risk parameters. Surprisingly, our study revealed that the prevalence of moderate and severe CKD, defined as eGFR <60 mL/min/1.73m², was 36.9% overall. The possible reasons for the high prevalence of CKD in our study are as follows. First, since the present study was performed at a hospital, it included many elderly patients, and thus the mean age was 66.3 years. The mean eGFR was 63.4 mL/min/1.73m² for all patients, 66.1 mL/min/1.73m² for men, and 60.9 mL/min/1.73m² for women. Therefore, the prevalence of CKD defined as eGFR<60 mL/min/1.73m² might have been high in our study. Second, GFR may be lower in Japanese than in Caucasians or African-Americans due to racial differences. Data from the U.S. NHANES III study indicated that the prevalence of CKD defined as eGFR<60 mL/min/1.73m² was 4.7% in the U.S. adult population aged 20 years or older, representing 8.3 million CKD patients (3). Most recently, the data from NIPPON DATA (National Integrated Project for Prospective Observation of Non-communicable Disease And its Trends in the Aged) 90 indicated that the prevalence of CKD defined as eGFR<60 mL/min/1.73m² was 6.7% among the 7,316 participants in a community population in Japan (23). Although our results are not directly comparable with these population-based studies, the prevalence of moderate and severe CKD might be higher in Japan compared to

the U.S. To confirm these relationships, a longitudinal study is needed.

Next, we demonstrated a stepwise increase in baPWV from CKD stages 1 to 5. Importantly, several studies across a broad spectrum of populations have shown that the risks of death and cardiovascular events are greatly increased in patients with an eGFR of less than 45 mL/min/1.73m² (2, 6). Therefore, we also investigated the findings when stage 3 was divided into two subsets: stage 3a, which included patients with an eGFR of 45 to 59; and stage 3b, which included those with an eGFR of 30 to 44 mL/min/1.73m². The baPWV of stage 3b patients was significantly higher than that of stage 3a patients, independent of age, systolic BP and other traditional risk factors. eGFR 45 mL/min/1.73m² may thus be a useful cut off value to predict arterial stiffness in CKD. These findings may explain, at least in part, why decreased eGFR is associated with greater risk for CVD and suggest that eGFR could be a useful surrogate end point for CVD.

The clinical significance of PWV is thought to relate not only to structural changes within the vascular wall but also to adverse hemodynamic effects (24, 25). These include an increase in systolic blood pressure and pulse pressure and, hence, an increase in dynamic left ventricular load. Meanwhile, decreased GFR may cause latent volume retention in patients with earlier stages of CKD. Furthermore, various factors that deteriorate vascular function, such as the renin-angiotensin-aldosterone system, homocysteine, oxidative stress, and inflammation, are also reportedly activated not only in patients with end-stage renal disease, but also those with earlier stages of CKD (1, 25-28). Anemia is also a key factor to determine the prognosis not only of dialysis patients and CKD but also of CVD. The published clinical and laboratory data suggest that anemia, CVD and CKD have a close

interrelationship, each worsening the others to result in a 'vicious cycle' of disease progression which Silberberg et al. have named the Cardio-Renal-Anemia Syndrome (29). There was a significant and negative correlation between baPWV and hemoglobin, although the statistical significance disappeared in the multiple regression analysis. Thus, decreased GFR may be associated with an increased level of traditional and nontraditional CVD risk factors.

Diabetic nephropathy is the leading cause of end-stage renal disease in the United States, Europe and Japan. Diabetes mellitus is associated with an increased risk of CVD (12). In the present study, eGFR was negatively associated with baPWV in both a subgroup with and one without diabetes. Meanwhile, a multiple regression analysis revealed that the correlation between baPWV and diabetes was statistically significant but not strong. Although the severity of diabetes may determine both eGFR and baPWV, eGFR seems to be a useful predictive marker of baPWV both in diabetes and non-diabetes.

Proteinuria is known as a risk factor for the development of end-stage renal disease and CVD. In the present study, the prevalence of proteinuria was positively correlated with the progression of CKD stage. The presence of proteinuria was associated with increased baPWV; however, the statistical significance disappeared after adjustment for traditional risk factors. Many factors thus seemed to play a role in the appearance of proteinuria and in its association with baPWV.

Our study had several limitations. First, the MDRD equation also includes age, sex and serum creatinine, and baPWV is itself closely associated with age. To minimize the influence of age and gender, we performed a multivariate analysis adjusted for age, gender,

and other conventional risk factors and showed that the independent contribution of eGFR to baPWV remained significant. Therefore, we believe that decreased eGFR was actually related to increased baPWV. Secondly, the MDRD equation for Japanese subjects, even after modification, has still not been completely approved for use. Thus, it is crucial that the new equation proposed by the Japanese Society of Nephrology be validated. Thirdly, we did not measure certain metabolic parameters in all subjects, such as triglycerides, HDL cholesterol, LDL cholesterol, uric acid, HbA1c and fasting glucose. We did not include these parameters in the multiple regression analysis, and thus we could not conclude that there was an interaction between these parameters and baPWV. Finally, the present study was a cross-sectional study and only included patients from a rural population aged 15 to 92 years who underwent medical examinations. Future longitudinal studies with a larger number of subjects from the general population will be needed to determine the association between risk factors for increased arterial stiffness and the progression of CKD.

In conclusion, our study suggests that the new eGFR calculated using the MDRD equation for Japanese patients is closely associated with an increase in arterial stiffness, and is a useful predictive marker for the development of atherosclerosis and CVD.

REFERENCES

1. Sarnak MJ, Levey AS, Schoolwerth AC, *et al*: American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation* 2003; **28**: 2154-2169.
2. Anavekar NS, McMurray JJ, Velazquez EJ, *et al*: Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med* 2004; **351**: 1285-1295.
3. Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS: Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2003; **41**: 1-12.
4. Mann JF, Gerstein HC, Yi QL, *et al*: Development of renal disease in people at high cardiovascular risk: results of the HOPE randomized study. *J Am Soc Nephrol* 2003; **14**: 641-647.
5. Muntner P, He J, Astor BC, Folsom AR, Coresh J: Traditional and nontraditional risk factors predict coronary heart disease in chronic kidney disease: results from the atherosclerosis risk in communities study. *J Am Soc Nephrol* 2005; **16**: 529-538.
6. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY: Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004; **351**: 1296-1305.

7. Levey AS, Beto JA, Coronado BE, *et al*: Controlling the epidemic of cardiovascular disease in chronic renal disease: what do we know? What do we need to learn? Where do we go from here? National Kidney Foundation Task Force on Cardiovascular Disease. *Am J Kidney Dis* 1998; **32**: 853-906.
8. National Kidney Foundation: K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis* 2002; **39**: Suppl 1:S1–S266
9. Lehmann ED: Clinical value of aortic pulse-wave velocity measurement. *Lancet* 1999; **354**: 528-529.
10. Asmar R, Benetos A, Topouchian J, *et al*: Assessment of arterial distensibility by automatic pulse wave velocity measurement. Validation and clinical application studies. *Hypertension* 1995; **26**: 485-490.
11. Blacher J, Guerin AP, Pannier B, Marchais SJ, Safar ME, London GM: Impact of aortic stiffness on survival in end-stage renal disease. *Circulation* 1999; **99**: 2434-2439.
12. Shoji T, Emoto M, Shinohara K, *et al*: Diabetes mellitus, aortic stiffness, and cardiovascular mortality in end-stage renal disease. *J Am Soc Nephrol* 2001; **12**: 2117-2124.
13. Munakata M, Nunokawa T, Yoshinaga K, Toyota T: Brachial-Ankle Pulse Wave Velocity Is an Independent Risk Factor for Microalbuminuria in Patients with Essential Hypertension—A Japanese Trial on the Prognostic Implication of Pulse Wave Velocity (J-TOPP). *Hypertens Res* 2006; **29**: 515-521.

14. Saijo Y, Yoshioka E, Fukui T, *et al*: Metabolic syndrome, C-reactive protein and increased arterial stiffness in Japanese subjects. *Hypertens Res* 2006; **29**: 589-596.
15. Wang MC, Tsai WC, Chen JY, Huang JJ: Stepwise increase in arterial stiffness corresponding with the stages of chronic kidney disease. *Am J Kidney Dis* 2005; **45**: 494-501.
16. Briet M, Bozec E, Laurent S, *et al*: Arterial stiffness and enlargement in mild-to-moderate chronic kidney disease. *Kidney Int* 2006; **69**: 350-357.
17. Ohya Y, Iseki K, Iseki C, *et al*: Increased pulse wave velocity is associated with low creatinine clearance and proteinuria in a screened cohort. *Am J Kidney Dis* 2006; **47**: 790-797.
18. Imai E, Horio M, Nitta K, *et al*: Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. *Clin Exp Nephrol* 2007; **11**: 41-50.
19. Horio M, Orita Y: Comparison of Jaffe rate assay and enzymatic method for the measurement of creatinine clearance. *Jpn J Nephrol* 1996; **38**: 296-299.
20. Coresh J, Astor BC, McQuillan G, *et al*: Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *Am J Kidney Dis* 2002; **39**: 920-929.
21. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999; **130**: 461-470.

22. Brosius FC 3rd, Hostetter TH, Kelepouris E, *et al*: Detection of chronic kidney disease in patients with or at increased risk of cardiovascular disease: a science advisory from the American Heart Association Kidney And Cardiovascular Disease Council; the Councils on High Blood Pressure Research, Cardiovascular Disease in the Young, and Epidemiology and Prevention; and the Quality of Care and Outcomes Research Interdisciplinary Working Group: developed in collaboration with the National Kidney Foundation. *Circulation* 2006; **114**: 1083-1087.
23. Nakamura K, Okamura T, Hayakawa T, *et al*: Chronic kidney disease is a risk factor for cardiovascular death in a community-based population in Japan: NIPPON DATA90. *Circ J* 2006; **70**: 954-959.
24. Yamashina A, Tomiyama H, Arai T, *et al*: Nomogram of the relation of brachial-ankle pulse wave velocity with blood pressure. *Hypertens Res* 2003; **26**: 801-806.
25. Safar ME, London GM, Plante GE: Arterial stiffness and kidney function. *Hypertension* 2004; **43**: 163-168.
26. London GM, Marchais SJ, Guerin AP, Metivier F, Adda H, Pannier B: Inflammation, arteriosclerosis, and cardiovascular therapy in hemodialysis patients. *Kidney Int Suppl* 2003; **84**: S88-93.
27. Tycho Vuurmans JL, Boer WH, Bos WJ, Blankestijn PJ, Koomans HA: Contribution of volume overload and angiotensin II to the increased pulse wave velocity of hemodialysis patients. *J Am Soc Nephrol* 2002; **13**: 177-183.
28. Fliser D, Kronenberg F, Kielstein JT, *et al*: Asymmetric dimethylarginine and progression of chronic kidney disease: the mild to moderate kidney disease study. *J Am*

Soc Nephrol 2005; **16**: 2456-2461.

29. Silverberg D: Outcomes of anaemia management in renal insufficiency and cardiac disease. *Nephrol Dial Transplant* 2003; **18** (Suppl 2): ii7-ii12.

Figure legends

Fig. 1. **Relationship between baPWV and eGFR in all 647 subjects.** GFR was estimated using the MDRD equation for Japanese proposed by the Japanese Society of Nephrology. baPWV was correlated significantly and negatively with eGFR ($r=-0.405$, $p<0.0001$). baPWV, brachial-ankle pulse wave velocity; eGFR, estimated glomerular filtration rate.


Fig. 2. **Distribution of eGFR and of CKD stages at baseline among the 647 subjects.** eGFR peaked in the range of 65-70 mL/min/1.73m² in men (black bar) and in the range of 60-65 mL/min/1.73m² in women (gray bar). The prevalences of CKD stages 1-5 were 2.6, 60.6, 34.2, 1.9 and 0.8%, representing 17, 392, 221, 12 and 5 patients, respectively. The prevalence of eGFR <60 mL/min/1.73m² was 36.9%, representing 238 patients (90 or 28.3% men and 148 or 45.0% women).

Fig. 3. **baPWV in patients with different stages of CKD.** (a) There was a stepwise increase in baPWVs, corresponding to advances of CKD through stages 1 to 5 ($p<0.0001$ for trend). (b) After adjustment for age, sex, systolic BP, BMI, HR, TC, prior CVD, smoking, medication for hypertension, medication for diabetes, and medication for dyslipidemia, significant differences in baPWV were observed across the different CKD stages ($p<0.0001$). When CKD stage 3, moderately reduced GFR, was divided at eGFR 45 mL/min/1.73m² (stage 3a: eGFR 45 to 59; stage 3b: eGFR 30 to 44 mL/min/1.73m²), (c) the adjusted baPWV in stage 3b was

found to be significantly greater than that in stage 3a ($p<0.0001$). The data are shown as the means \pm SD.


Fig. 4. **Score of proteinuria in patients with different stages of CKD.** There was a stepwise increase in the score of proteinuria, corresponding to advances in CKD through stages 1 to 5 ($p<0.0001$ for trend). The data are shown as the means \pm SD.

Figure 1




Nakagawa, N., *et al*

Figure 2




Nakagawa, N., *et al*

Figure 3A




Nakagawa, N., et al

Figure 3B




Nakagawa, N., *et al*

Figure 3C



Nakagawa, N., *et al*

Figure 4



Nakagawa, N., et al